



³Michigan State University, College of Osteopathic Medicine, East Lansing, MI, U.S.A

1. Morgan MD, Salmon LJ, Waller A, Roe JP, Pinczewski LA. Fifteen-year survival of endoscopic anterior cruciate ligament reconstruction in patients aged 18 years and younger. *Am J Sports Med.* 2016;44(2):384-392.
2. Fauno P, Rahr-Wagner L, Lind M. Risk for revision after anterior cruciate ligament reconstruction is higher among adolescents: results from the danish registry of knee ligament reconstruction. *Orthop J Sports Med.* 2014;2(10).
3. Brophy RH, Schmitz L, Wright RW, et al. Return to play and future ACL injury risk after ACL reconstruction in soccer athletes from the multicenter orthopaedic outcomes network (MOON) group. *Am J Sports Med.* 2012;40(11):2517-2522.
4. McCullough KA, Phelps KD, Spindler KP, et al. Return to high school- and college-level football after anterior cruciate ligament reconstruction: a Multicenter Orthopaedic Outcomes Network (MOON) cohort study. *Am J Sports Med.* 2012;40(11):2523-2529.
5. Shelbourne KD, Benner RW, Gray T. Return to sports and subsequent injury rates after revision anterior cruciate ligament reconstruction with patellar tendon autograft. *Am J Sports Med.* 2014;42(6):1395-1400.
6. WHO | Adolescent development. WHO. http://www.who.int/maternal_child_adolescent/topics/adolescence/dev/en/. Accessed February 12, 2016.
7. Rice SG, American Academy of Pediatrics Council on Sports Medicine and Fitness. Medical conditions affecting sports participation. *Pediatrics.* 2008;121(4):841-848.